arXiv:1407.4326v1 [math.GR] 16 Jul 2014

ON DIVISIBILITY GRAPH FOR SIMPLE ZASSENHAUS GROUPS

A. ABDOLGHAFOURIAN AND MOHAMMAD A. IRANMANESH

Abstract. The divisibility graph D(G) for a finite group G is a graph with vertex set cs(G)$\setminus\{1\}$ where cs(G) is the set of conjugacy class sizes of G. Two vertices a and b are adjacent whenever a divides b or b divides a. In this paper we will find D(G) where G is a simple Zassenhaus group.

1. Introduction and Prelimits

There are several graphs associated to algebraic structures, specially finite groups, and many interesting results have been obtained recently (see for example [4, 8, 9, 10]). In [3] a new graph namely *divisibility graph* which is related to a set of positive integers have been introduced. The divisibility graph, $\overrightarrow{D}(X)$ is a graph with vertex set $X^* = X \setminus \{1\}$ and there is an arc between two vertices a and b if and only if a divides b. Since this graph is never strongly connected, we consider its underlying graph D(X) without changing the name for convenience. Also we use D(G) instead of D(cs(G)) where cs(G) is the set of conjugacy class sizes of elements of G.

All groups considered here are finite. Let H be a subgroup of a group G. We use $[G : H] = \{Hg|\ g \in G\}$ to denote the right cosets of H in G. The conjugacy class of an element $g \in G$, will denote by g^G which is the set of all $x^{-1}gx$ where $x \in G$. Also we use cs(G) for the set of conjugacy class sizes of elements of G. The conjugate of $h \in H$ by $g \in G$ and the conjugate of H by g are $h^g = g^{-1}hg$ and $H^g = g^{-1}Hg$ respectively. Also by $C_G(g) = \{x \in G|\ g^x = g\ \}$, $Z(H) = \{z \in$

2010 *Mathematics Subject Classification.* 20D05, 05C25.
Key words and phrases. Divisibility graph, Zassenhaus groups, Simple groups.
Corresponding author: Mohammad A. Iranmanesh.

$G|\ g^z = g\ for\ every\ g \in G\ \}$ and $N_G(H) = \{g \in G|\ H^g = g\ \}$, we denote the centralizer of g in G, the center of G and the normalizer of H in G respectively. An involution of G is an element of order 2. By H is disjoint from its conjugates we mean that $H \cap H^g = \{e\}$ *or* H, for every $g \in G$. We write $G =< g >$, to show that G is generated by g. For a finite set X we use $|X|$ for cardinality of X and $\gcd(a, b)$ is the greatest common divisor of a and b where a and b are positive integers. For undefined terminology and notation about groups we refer to [11].

Throughout the paper all graphs are finite and simple. Let Γ_1, Γ_2 be two graphs then $\Gamma = \Gamma_1 + \Gamma_2$ means Γ is disjoint union of Γ_1 and Γ_2. The complete graph with n vertices will denote by $\mathrm{K_n}$. For other notation about graphs we refer to [14].

The divisibility graph D(G), where G is a symmetric group or an alternating group studied in [1]. It is shown there that $\mathrm{D(S_n)}$ has at most two connected components and if it is not connected then one of its connected components is $\mathrm{K_1}$. Also we have proved that $\mathrm{D(A_n)}$ has at most three connected components and if it is not connected then two of its connected components are $\mathrm{K_1}$.

In this paper we consider D(G) where G is a simple Zassenhaus group.

Definition 1. [7, Chapter 13] *A permutation group G is a Zassenhaus group if it is doubly transitive in which only the identity fixes three letters.*

Theorem 1. [7, Chapter 13] *Let G be a Zassenhaus group of degree $n + 1$ and let N be the subgroup fixing a letter. Then we have*

(i) *N is a Frobenius group with kernel K of order n and complement H.*

(ii) *K is a Hall subgroup of G, K is disjoint from its conjugates, $C_G(k_0) \leq K$ for every $k_0 \in K$ and $N = N_G(K)$.*

(iii) *H is a subgroup of G fixing two letters, H is disjoint from its conjugates in G, and $[N_G(H) : H] = 2$.*

(iv) *$|G| = en(n + 1)$, where $e = |H|$ and e divides $n - 1$.*

(v) *If e is even then K is abelian.*

(vi) *If $e \geq (n-1)/2$ then K is an elementary abelian p-group for some prime p.*

(vii) *If G is simple and e is odd then H is cyclic. Also G has only one conjugacy class of involutions of size $e(n+1)$ if n is even and of size en if n is odd.*

In this case we say that G is of type (H, K).

Also G has only one conjugacy class of involutions of size $e(n+1)$ if n is even and of size en if n is odd.

In this case we say that G is of type (H, K). As it is stated in [7, Chapter 16, Page 488], PSL(2, q), the projective special linear group of dimension two over a finite field $\mathbb{F}_q$ of order $q > 3$, and Sz(q), Suzuki simple groups of order $q^2(q-1)(q^2+1)$ where $q = 2^m$ and $m > 1$ is an odd number are the only simple Zassenhaus groups. (For more details about the structure of PSL(2, q) and Sz(q) we refer to [5] and [12] respectively.)

According to the notations of Theorem 1, the projective special linear groups PSL(2, q) are Zassenhaus group of type (H, K) of degree $n+1$ where $|K| = n = q$, $e = (n-1)/2$ if n is odd, and $e = n-1$ otherwise. Also H is cyclic group that inverted by an involution [7, Chapter 13]. So by Theorem 1, $|G| = q(q^2-1)/2$ if q is odd and $|G| = q(q^2-1)$ otherwise. It is proved in [5, Chapter XII.] that the projective special linear groups, PSL(2, q), has a cyclic subgroup namely L of order $(q+1)/\gcd(2, q+1)$ such that $x^{\mathrm{PSL(2,q)}} \cap L = \{x, x^{-1}\}$ for every $x \in L$. L is disjoint from its conjugates in G and $[N_G(L) : L] = 2$.

Also by Theorem 1, we see that the Suzuki simple groups, Sz(q), are Zassenhaus group of type (H, K) with H is cyclic of order $q-1$ and K is a nonabelian 2-group of order q^2 with elementary abelian center of order q and $N = HK$ is the subgroup fixing a letter [7, Page 466]. So by Theorem 1, $|G| = q^2(q-1)(q^2+1)$ where $q = 2^m$ and $m > 1$ is an odd number.

Not that by [6] the Suzuki simple groups, Sz(q), can be defined as a subgroup of the group $\mathrm{SL}_4(q)$. In this representation, $K =< (\alpha,\beta) \mid \alpha,\ \beta \in \mathbb{F}_q >$ and $Z(K) = \{(0,\beta) \mid \beta \in \mathbb{F}_q\}$ where (α,β) is defined

$$(\alpha,\beta) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ \alpha & 1 & 0 & 0 \\ \alpha^{1+r}+\beta & \alpha^r & 1 & 0 \\ \alpha^{2+r}+\alpha\beta+\beta^r & \beta & \alpha & 1 \end{bmatrix},$$

with multiplicity $(\alpha,\beta)(\gamma,\delta) = (\alpha+\gamma, \alpha\gamma^r+\beta+\delta)$ and $r^2 = 2q$. Note that this implies that all $1 \neq x \in Z(K)$ are involutions.

The following three lemmas about Suzuki groups will be useful.

Lemma 2. [12, Proposition 16] Sz(q) *contains abelian subgroups* $A_0,\ A_1$ *and* A_2 *of order* $q-1$ *and* $q+r+1$ *and* $q-r+1,\ (r^2=2q)$, *respectively.* $C_G(x) = A_i$ *for every* $x \in A_i$ *and* $i = 0,1,2$. *Moreover* $[N_G(A_i) : A_i] = 4$ *for* $i = 1,2$.

Lemma 3. [6, Lemma 3.3] *The subgroups* A_i*'s are disjoint from their conjugates for* $i = 0,\ 1,\ 2$.

By Theorem 1, we see that $H = A_0$.

Lemma 4. [12, Lemma 3.2., Part (h)] *Let* $x = (\alpha,\beta) \in K$, *where* $\alpha \neq 0$. *Then* $C_G(x) =< (\alpha,0), Z(K) >$, *where* $Z(K) = \{(0,\xi) : \xi \in \mathbb{F}_q\}$.

Now we prove the following lemma.

Lemma 5. *Let* G *be a group and* H *a subgroup of* G *which is disjoint from its conjugates in* G. *Suppose* h *is a non identity element of* H, $N_G(H) = N$, *and* $C_G(h) = C$. *Then* $|N| = |h^G \cap H||C|$.

Proof. Let $c \in C$. So $h^c = h$ and $H^c \cap H \neq \{e\}$. Since H is disjoint from its conjugates, we can conclude $H^c = H$. Hence $c \in N$ and $C \leq N$. Define a map θ from $[N : C]$ to $h^G \cap H$, by $\theta(Cg) = h^g$. Suppose g_0 is an arbitrary element of G. Since H is disjoint from its conjugates, we conclude $h^{g_0} \in H$ if and only if $g_0 \in N$.

Also $Cx = Cy$ for $x, y \in N$ if and only if $xy^{-1} \in C$, which implies that θ is well defined and injective. It is easy to see that θ is also surjective. This completes the proof. □

In the next section we will calculate the conjugacy class sizes of elements of the projective special linear groups PSL(2, q) and the Suzuki groups Sz(q), when they are simple. The structure of divisibility graph for these groups will also investigate.

2. The structure of divisibility graph for simple Zassenhaus groups

This section includes two theorems. We will determine the structure of divisibility graphs for PSL(2, q) and Sz(q) when they are simple.

Theorem 6. *Let* $G = \mathrm{PSL}(2, \mathrm{q})$. *Then* D(G) *is one of the graphs of the following list:*

(i) $3\mathrm{K}_1$

(ii) $\mathrm{K}_2 + 2\mathrm{K}_1$.

Proof. To calculate the conjugacy class sizes of non identity elements of G, we consider three subgroups H, K and L which their orders are pairwise coprime. So the non identity elements of these groups are not conjugate.

First consider the subgroup H. Let $H =< h_0 >$. By the structure of G, we know that H is inverted by an involution. So every elements of H is cinjugate to its inverse and $N_G(H) \neq C_G(H)$. Also $H \leq C_G(h_0) = C_G(H)$. Since by part (iii) of Theorem 1, $[N_G(H) : H] = 2$, then $C_G(h_0) = H$. This implies $|h_0^G| = q(q+1)$.

Consider an arbitrary element $1 \neq h \in H$. Clearly $C_G(h_0) \leq C_G(h)$. If $C_G(h_0) = C_G(h)$ then $|h^G| = q(q+1)$ and h is different from its inverse as h_0 is. If $C_G(h_0) \neq C_G(h)$ then, since H is disjoint from its conjugate, $C_G(h) = N_G(H)$. So by Lemma 5, h has only one conjugate in H. Hence $h = h^{-1}$. This means that h is an involution and $|h^G| = q(q+1)/2$. Therefore if e is even, there exist one conjugacy classes of size $q(q+1)/2$ related to $h_0^{e/2}$ and $(e-2)/2$ conjugacy class of

size $q(q+1)$ in G, related to other non identity elements of H. If e is odd then G has $(e-1)/2$ conjugacy classes of size $q(q+1)$ that contains non identity elements of H.

Now consider the subgroup K. First suppose $q = 2^\alpha$. In this case by Theorem 1 parts (vi) and (vii), K is an elementary abelian 2-group and G has only one conjugacy class of involutions. So every element of K contains in the unique conjugacy class of involutions of size (q^2-1).

Suppose $q = p^\alpha$ is an odd number. By parts (ii) and (vi) of Theorem 1, $K = C_G(y)$ for each $y \in K$. Since $[N : K] = e$, by Lemma 5, $|y^G \cap K| = e$. So we can partition $K \setminus \{1\}$ into two disjoint subset of length e, each of them contains conjugate elements. Hence in this case, there are two conjugacy classes in G of size $(q^2-1)/2$ that contains elements of K.

Finally consider the subgroup L. Since for every $x \in L$ we have $x^{\mathrm{PSL(2,q)}} \cap L = \{x, x^{-1}\}$ by Lemma 5, if $x \in L$ is not an involution then we have $|C_G(x)| = |N_G(L)|/2 = |L|$ and if $x \in L$ is an involution then $|C_G(x)| = |N_G(L)| = 2|L|$. So $|L|$ is even then there exist $(|L|-2)/2$ conjugacy classes in G of size $\gcd(2, q+1)eq$ and one conjugacy class of size $\gcd(2, q+1)eq/2$ that contain elements of L. If $|L|$ is odd, then there exist $(|L|-1)/2$ conjugacy classes in G of size $\gcd(2, q+1)eq$ that contain elements of L.

We must consider two following cases:

(Case 1) $G = \mathrm{PSL}(2, 2^{\mathrm{k}})$. In this case $e = n-1 = q-1$ is an odd number. By using class equation of G we have;

$$1 \ + \ (|L|-1)/2 \ . \ eq \ + \ (q-1)(q+1) \ + \ ((e-1)/2) \ . \ q(q+1) \ =$$
$$1 \ + \ q^2(q-1)/2 \ + \ (q^2-1) \ + \ (q-2)q(q+1)/2 \ = \ |G|.$$

(Case 2) $G = \mathrm{PSL}(2, \mathrm{p^k})$ where p is an odd prime. In this case $e = (n-1)/2 = (q-1)/2$. First assume that e is odd. In this case we have;

$$1 \ + \ ((|L|-2) \ / \ 2) \ . \ 2eq \ + \ eq \ + \ 2 \ . \ e(q+1) \ + \ ((e-1)/2) \ . \ q(q+1) \ =$$

$$1 \ + q(q-1)(q-3)/4 \ + \ q(q-1)/2 \ + \ (q^2-1) \ + \ (q-3)q(q+1)/4 \ = |G|.$$

In case where e is even we have;

$$1 \ + \ ((|L|-1)/2) \ . \ 2eq \ + \ 2 \ .e(q+1) \ + \ ((e-2)/2) \ . \ q(q+1) \ + \ q(q+1)/2 =$$

$$1 \ + \ q(q-1)^2/4 \ + \ (q^2-1) \ + \ (q-5)q(q+1)/4 \ + \ q(q+1)/2 = |G|.$$

Hence by calculating all conjugacy classes of $\mathrm{PSL}(2, \mathrm{q})$, we see that $\mathrm{D}(\mathrm{PSL}(2, \mathrm{q}))$ is $3\mathrm{K}_1$ or $\mathrm{K}_2 + 2\mathrm{K}_1$. (See Figure 1.) □



FIGURE 1. The graph $\mathrm{D}(\mathrm{PSL}(2, \mathrm{q}))$

Now we calculate the conjugacy class sizes of Suzuki simple groups.

Theorem 7. *Let* $G = \mathrm{Sz(q)}$. *Then* $\mathrm{D(G)} = \mathrm{K}_2 + 3\mathrm{K}_1$.

Proof. By Lemmas 2 and 4, G has four subgroups $H,\ K,\ A_1$ and A_2 with pairwise coprime orders.

We start by calculating the conjugacy class sizes of elements of H. By Lemma 2, $C_G(h) = H$ for every $1 \neq h \in H$. Since H is disjoint from its conjugates, by Lemma 5, h has $[N_G(H) : C_G(h)] = 2$ conjugates in H. So there are $(q-2)/2$ conjugacy classes in G of size $q^2(q^2+1)$ which contains elements of H.

Consider the conjugacy classes of elements of K. Let $x_0 = (\alpha, \beta) \in K$ be an arbitrary element. If $1 \neq x_0 \in Z(K)$ then x_0 is an involution and by Theorem 1 part

(vii), G has only one class of involutions of size $(q-1)(q^2+1)$. If $x_0 \in K \setminus Z(K)$ then by Lemma 4, $C_G(x_0) = \{(x,y)|\ y \in \mathbb{F}_q,\ x = 0\ or\ \alpha\}$. This means that $|C_G(x_0)| = 2q$ and $|x_0^G| = q(q-1)(q^2+1)/2$. Since K is disjoint from its conjugates, by Lemma 5, x_0 has $[K : C_G(x_0)] = q(q-1)/2$ conjugates in $K \setminus Z(K)$. So there exist $(q^2-q)/(q(q-1)/2) = 2$ conjugacy classes of size $q(q-1)(q^2+1)/2$ in G which contains elements of $K \setminus Z(K)$.

Finally we consider two subgroups A_1 and A_2. By Lemma 2, A_1 and A_2 are of order $n_1 = q+r+1$ and $n_2 = q-r+1$ respectively. Also $[N_G(A_i) : A_i] = 4$ and for every $x \in A_i$, $C_G(x) = A_i$ where $i = 1, 2$. So for every $x \in A_i,\ i = 1, 2$, we have $[N_G(A_i) : C_G(x)] = 4$ and by Lemmas 3 and 5, there are just four elements of A_i which are conjugate with x in G. Hence there are $(q+r)/4$ conjugacy classes of size $q^2(q-1)(q^2+1)/(q+r+1) = q^2(q-1)(q-r+1)$ which contains elements of A_1 and $(q-r)/4$ conjugacy classes of size $q^2(q-1)(q^2+1)/(q-r+1) = q^2(q-1)(q+r+1)$ which contains elements of A_2.

By using the class equation for G we see that

$$1 \ + \ (q-2)\ .\ q^2(q^2+1)/2 \ + \ (q-1)(q^2+1) \ + \ 2\ .\ q(q-1)(q^2+1)/2 +$$

$$(q+r)\ .\ q^2(q-1)(q-r+1)/4 \ + \ (q-r)\ .\ q^2(q-1)(q+r+1)/4 = |G|.$$

This implies that D(Sz(q)) = $K_2 + 3K_1$. (See Figure 2.) □



FIGURE 2. The graph D(Sz(q))

REFERENCES

[1] Abdolghafourian A and Iranmanesh M A, Divisibility graph for symmetric and alternating groups. *Accepted*

[2] Bertram E A, Herzog M and Mann A, On a graph related to conjugacy classes of groups. *Bull. London Math. Soc.* **22** (1990) 569–575

[3] Camina A R and Camina R D, The influence of conjugacy class sizes on the structure of finite groups: a survey, *Asian-Eur. J. Math.* **4** (2011) 559–588

[4] Darafsheh M R and Assari A, Normal edge-transitive Cayley graphs on non-abelian groups of order 4p, where p is a prime number, *Sci China Ser A* **56** (2013) 213–219

[5] Dickson L E, Linear groups, with an Exposition of the Galois Field Theory. (1901) (Teubner Leipzig)

[6] Fang X G and Praeger C E, Finite two arc transitive graphs admitting a Suzuki simple group, *Comm. Algeb.* **27** (1999) 3727–3754

[7] Gorenstein D, Finite Groups, 2nd Edittion, (1980) (Chelsea, New York)

[8] Hafezieh R and Iranmanesh M A, Bipartite divisor graph for the product of subsets of integers, *Bull. of the Aust. Math. Soc.* **87** (2013) 288–297

[9] Iranmanesh M A and Praeger C E, Bipartite divisor graphs for integer subsets, *Graphs Combin.* **26** (2010) 95–105

[10] Karla H and Gumber D, A note on conjugacy classes of finite groups, *Proc. Indian Acad. Sci. (Math. Sci.)* **124** (2014) 31–36

[11] Rose J S, A course ongroup theory, (1978) (Cambridge University Press)

[12] Suzuki M, On a class of doubly transitive groups. *Annal. Math.* **75** (1962) 105–145

[13] Ward H N, On Ree's series of simple groups, *Trans. Amer. Math. Soc.* **121** (1966) 62–89

[14] West D B, Introduction to Graph Theory, 2nd ed, (2000) (Englewood Cliffs, NJ: Prentice-Hall)

A. Abdolghafourian, Department of Mathematics

Yazd University, Yazd, 89195-741, Iran

E-mail address: Adeleh Abdolghafourian@stu.yazd.ac.ir

M. A. Iranmanesh, Department of Mathematics

Yazd University, Yazd, 89195-741 , Iran

E-mail address: iranmanesh@yazd.ac.ir